

The detection and determination of diene hydrocarbons with a conjugated system of double bonds. I. V. I. Kraslov. J. Applied Chem. (U. S. S. R.) 14, 101-173 (1941); translated in *Foreign Petroleum Tech.* 9, 344-363 (1941). The Kunkmann-Kaufmann method (L. A. 30, 7886) cannot be used universally for the detn. of diene hydrocarbons with conjugated double bonds. The excessively low values of the Kaufmann diene numbers are due to side reactions of the polymerization of dienes. Diene hydrocarbons are distinguished from monoolefins by the generation of considerable amounts of HBr when brominated in a soln. of CCl_4 . The Mellinley method is suitable for detection of dienes alone or in mixts. of various hydrocarbons. II. V. I. Kraslov and A. V. Shpash. J. Applied Chem. (U. S. S. R.) 14, 104-111 (in German, 1941); translated in *Foreign Petroleum Tech.* 9, 360-361 (1941).—Bromination in CCl_4 is specific for diene hydrocarbons with conjugated double bonds and is accompanied by the generation of considerable amounts of halogen acids. Hydrocarbon chains with separated double bonds are brominated similarly to olefins, with an insignificant loss of halogen acid. The sensitivity limits of the bromination reaction for mixts. of olefins with 1,3-dienes is 1% of the latter, while for mixts. of said dienes with dienes this limit is at 0.5% of dienes. The experiments were carried out with linear oil and mixts. of cyclohexene and decalin, and with ketone and dicarbenes, by the Mellinley method of degt., 1 nos. A. A. Boehmlich

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0"

Characterization of α , β -unsaturated ketones. II. V. F. Kaufer and J. I. Rakicher, *J. Gen. Chem. (U.S. S. R.)* 13, No. 9-13 (1933) (English summary); cf. *C. A.* 29, 3188. It was shown that the displacement of ketones by aldehydes from α , β -unsubstituted ketones is connected with the preliminary hydrolytic cleavage of the latter into its components. Buff., treated with methyl oxide in the presence of dil. NaOH, yielded small amounts of benzylidenemalonic and benzylideneacetoxy oxides, the latter being isolated as the tetrabromide, m. 118° (from EtOH). The reactions were conducted by prolonged standing at about 10°, with a longer period noticeably increasing the yield of benzylidenacetone. Me₂CO was shown to be inert to condensations on treatment with piperidine acetate or prolonged heating on a steam bath; the same result was obtained when MeEt₂CO was used. However, 25 g. Buff., 30 g. Me₂CO, 5 g. piperidine and 5 g. AcOH yielded, after 20 hrs. heating, 71% benzylidenacetone, m. 141.8°, while similar reaction with MeEt₂CO gave crude benzylideneacetoxy oxide (m. 173°), identified as the tetrabromide. Thus, piperidine acetate may be recommended for aldehyde-ketone condensations. G. M. Karasikoff

G. M. Kondajing

APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0"

Preparation of enanthylidenacetone. V. I. B. 1943
I. P. Vladimirov, M. S. Kazanitina, Z. S. Ullila, Z. B.
Prokof'eva and I. I. Radzhev, *J. Gen. Chem. (U. S. S. R.)* 13,
814-17 (1943) (English summary).—It was shown that the
product obtained by Kupe and Hünigtsch (C. A. J.,
1938) from condensation of Me₂CO with enanthio in the
cold in the presence of 1% NaCl is not enanthyliden-
acetone but decan-4-ol-*o*-one. Authentic enanthyliden-
acetone was prepared, as follows: 3 g. naphthalene and 3 g. glu-
ceral AcOH, in 50 g. dry Me₂CO, were treated dropwise with
29 g. enanthio while refluxing the tubat. on a steam bath; after the addition, which requires 4 hrs., the mixt. was
heated for 8 hrs., cooled, the eq. layer sept., while the org.
layer was freed of Me₂CO, dried with Et₂O, dried over
Na₂SO₄, and distilled, to yield enanthylidenacetone, bp 103-
104°, n_D²⁰ 1.4480, d₄²⁰ 0.8473; yield, 16-17 g. The product
yields the normal positive iodform reaction. The product
obtained by K. and H. yields enanthylidenacetone after
dehydration over KHSO₄-Na₂SO₄, by repeated distn. at 32
mm.

10

450-51A METALLURGICAL LITERATURE CLASSIFICATION												450-51A METALLURGICAL LITERATURE CLASSIFICATION																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
FROM 1910-1930												TO 1940-1960																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
SEARCHED MAY 1970												SEARCHED MAY 1970																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
160000	40	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000	1001	1002	1003	1004	1005	1006	1007	1008	1009	10010	10011	10012	10013	10014	10015	10016	10017	10018	10019	10020	10021	10022	10023	10024	10025	10026	10027	10028	10029	10030	10031	10032	10033	10034	10035	10036	10037	10038	10039	10040	10041	10042	10043	10044	10045	10046	10047	10048	10049	10050	10051	10052	10053	10054	10055	10056	10057	10058	10059	10060	10061	10062	10063	10064	10065	10066	10067	10068	10069	10070	10071	10072	10073	10074	10075	10076	10077	10078	10079	10080	10081	10082	10083	10084	10085	10086	10087	10088	10089	10090	10091	10092	10093	10094	10095	10096	10097	10098	10099	100100	100101	100102	100103	100104	100105	100106	100107	100108	100109	100110	100111	100112	100113	100114	100115	100116	100117	100118	100119	100120	100121	100122	100123	100124	100125	100126	100127	100128	100129	100130	100131	100132	100133	100134	100135	100136	100137	100138	100139	100140	100141	100142	100143	100144	100145	100146	100147	100148	100149	100150	100151	100152	100153	100154	100155	100156	100157	100158	100159	100160	100161	100162	100163	100164	100165	100166	100167	100168	100169	100170	100171	100172	100173	100174	100175	100176	100177	100178	100179	100180	100181	100182	100183	100184	100185	100186	100187	100188	100189	100190	100191	100192	100193	100194	100195	100196	100197	100198	100199	100200	100201	100202	100203	100204	100205	100206	100207	100208	100209	100210	100211	100212	100213	100214	100215	100216	100217	100218	100219	100220	100221	100222	100223	100224	100225	100226	100227	100228	100229	100230	100231	100232	100233	100234	100235	100236	100237	100238	100239	100240	100241	100242	100243	100244	100245	100246	100247	100248	100249	100250	100251	100252	100253	100254	100255	100256	100257	100258	100259	100260	100261	100262	100263	100264	100265	100266	100267	100268	100269	100270	100271	100272	100273	100274	100275	100276	100277	100278	100279	100280	100281	100282	100283	100284	100285	100286	100287	100288	100289	100290	100291	100292	100293	100294	100295	100296	100297	100298	100299	100300	100301	100302	100303	100304	100305	100306	100307	100308	100309	100310	100311	100312	100313	100314	100315	100316	100317	100318	100319	100320	100321	100322	100323	100324	100325	100326	100327	100328	100329	100330	100331	100332	100333	100334	100335	100336	100337	100338	100339	100340	100341	100342	100343	100344	100345	100346	100347	100348	100349	100350	100351	100352	100353	100354	100355	100356	100357	100358	100359	100360	100361	100362	100363	100364	100365	100366	100367	100368	100369	100370	100371	100372	100373	100374	100375	100376	100377	100378	100379	100380	100381	100382	100383	100384	100385	100386	100387	100388	100389	100390	100391	100392	100393	100394	100395	100396	100397	100398	100399	100400	100401	100402	100403</td

PROCESSED AND PREDICTED 4084

10

2d.


Synthesis of new hydrocarbons with a quadrangular system of double bonds. IV. V. A. Egorov, I. F. Vilkovskaya, M. S. Kozlikhina and I. I. Valkher. J. Gen. Chem. (U. S. S. R.) 13, 818-23 (1943) (English summary); cf. C. A. 39, 4739 and following above.—Methyl (from 14.8 g. MeI) was added dropwise over 5 hrs. to 13.4 g. malonitrile/diisopropylacetone in Et₂O at -15°; after standing overnight and decomps. by ice and NH₄Cl, there was obtained 18% 2-methyl-1,3-dodeadiene, bp 88°, η_2° 1.4492, d_2^{20} 0.7718. The product, heated for 20 hrs. in a steam bath in a sealed tube with malonic anhydride in toluene, yielded an adduct, apparently of the trimer, which could not be reduced to the cryst. state. Similar reaction using Ethigell gave a mixt. of 3-methyl-2,4-hexadecadiene and 3-alkyl-1,3-dodeadiene anhydride gives a mixt. of adducts of the monomer and the dimer of the hydrocarbon, in the form of a heavy oil. A similar prepn. using iso-Armidgill gave a mixt. of 2-isooxy-1,3-dodeadiene and 2,6-dimethyl-4,6-octadecadienoate, bp 120-34°, η_2° 1.4164, d_2^{20} 0.7806. G. M. Kosolapoff

AB-51A METALLURGICAL LITERATURE CLASSIFICATION

SEARCHED INDEXED 483040 M17 OME 101

140082 94

REF ID	TITLE	SUBJ	EFFECTIVENESS												TECHN. ADAPTABILITY											
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27




Ca


10

Effecting the Grignard reaction with α,β -unsaturated ketones. V. V. I. Ruzav. *J. Gen. Chem. (U.S.S.R.)* 14, 84-7(1944) (English summary); cf. preceding chart. Simultaneous addn. of EtMgBr (in Et₂O) and methyl oxide (in Et₂O) to BaO kept at -10°, followed by the usual work-up and dehydration of the crude tertiary alk. with CaH(CO)₂O, gave an av. yield of 50.9% 2,4-dimethyl-2,4-hexadiene, b. 114-15°, thus showing that, in reactions of Grignard reagents with α,β -unsatd. ketones, the diln. and employment of low temp. favors the normal reaction course. The same reaction, conducted at 15°, gave 17.8% of crude octadienes, consisting of 2-methyl-4-methyl-1,3-pentadiene, b. 111-12°, d₄²⁰ 0.7633, n_D²⁰ 1.4424, and an undetd. amt. of the higher-boiling isomer. The product of the -10° reaction was sepd., in the ratio of 1:1, into the above pentadiene and the hexadiene, b. 114-15°, d₄²⁰ 0.7642, n_D²⁰ 1.4430. The structures of the compds. were dtd. by oxidation. G. M. Kosolapoff

ATA-SEA METALLURGICAL LITERATURE CLASSIFICATION



		10-42-102-000179		PROCESSES AND PRODUCTS OF SIDER			
		<p><i>Halogenation of unsaturated hydrocarbons. IV.</i></p> <p>I. Basov (Sverdlovsk State Univ.). <i>J. Gen. Chem. (U.S.S.R.)</i> 13, 1001-6 (1943); cf. <i>C.A.</i> 39, 4704.—The Macallum method of iodine no. detn. is recommended for detection of structures at the quaternary C atom having a double bond, such as in olefins or in dienes. The amt. of such hydrocarbons in liquid-phase cracking products is shown to increase with rising b.p. of the fractions. A hypothesis for halogenation and hypohalogenation of olefins is proposed according to which the halogenating mol. is first fixed to the double bond, then the atomic bonds are cleaved, and the halogens are transformed into <i>statox nascondi</i>; this explains the possibility and the high reaction rate of such halogens with other materials present in the reaction mixt. Isoamylene (1.00 g.) and 8.9992 g. kerosene fraction with iodine no. 0.0 were mixed and 4.7063 g. of the <i>redu</i>. was treated for 0.5 hr. with 2.5 l. freshly prep'd. Cl water, and the org. layer was sepd., washed with dil. soda soln., dried, and analyzed for Cl and iodine; 19% of the isoamylene did not enter into reaction, thus showing direct chlorination of isoamylene; extension of the reaction to 1 hr. gave 17% unreacted isoamylene. The supposition that hypochlorination is accompanied by direct chlorination is also supported by the higher Cl content than calcd. for an HOCl addn. product. Combination of 2,4-dimethyl-2,4-hexadiene in CCl_4 at 0° for 0.5-2.0 hrs. showed that 8.8-20.2% of total utilized Br is used for transformation into HBr. G. V. Konopljoff</p>		10			
		450-514 METALLURGICAL LITERATURE CLASSIFICATION		UNITED STATES			
		SCIENTIFIC		INDUSTRIAL			
		SCIENCE & TECHNOLOGY	INDUSTRY & PRODUCTION	INDUSTRY & PRODUCTION			
		GENERAL	GENERAL	GENERAL			
		PHYSICS	CHEMISTRY	CHEMISTRY			
		MATHEMATICS	MATHEMATICS	MATHEMATICS			
		PHYSICAL CHEMISTRY	PHYSICAL CHEMISTRY	PHYSICAL CHEMISTRY			
		ORGANIC CHEMISTRY	ORGANIC CHEMISTRY	ORGANIC CHEMISTRY			
		INORGANIC CHEMISTRY	INORGANIC CHEMISTRY	INORGANIC CHEMISTRY			
		PETROCHEMISTRY	PETROCHEMISTRY	PETROCHEMISTRY			
		PETROLEUM	PETROLEUM	PETROLEUM			
		PETROGRAPHY	PETROGRAPHY	PETROGRAPHY			
		PETROPHYSICS	PETROPHYSICS	PETROPHYSICS			
		PETROPHYSICO-CHEMICAL METHODS	PETROPHYSICO-CHEMICAL METHODS	PETROPHYSICO-CHEMICAL METHODS			
		PETROPHYSICO-MATHEMATICAL METHODS	PETROPHYSICO-MATHEMATICAL METHODS	PETROPHYSICO-MATHEMATICAL METHODS			
		PETROPHYSICO-TECHNICAL METHODS	PETROPHYSICO-TECHNICAL METHODS	PETROPHYSICO-TECHNICAL METHODS			
		PETROPHYSICO-TECHNICAL EQUIPMENT	PETROPHYSICO-TECHNICAL EQUIPMENT	PETROPHYSICO-TECHNICAL EQUIPMENT			
		PETROPHYSICO-TECHNICAL INSTRUMENTS	PETROPHYSICO-TECHNICAL INSTRUMENTS	PETROPHYSICO-TECHNICAL INSTRUMENTS			
		PETROPHYSICO-TECHNICAL APPARATUS	PETROPHYSICO-TECHNICAL APPARATUS	PETROPHYSICO-TECHNICAL APPARATUS			
		PETROPHYSICO-TECHNICAL MACHINES	PETROPHYSICO-TECHNICAL MACHINES	PETROPHYSICO-TECHNICAL MACHINES			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY	PETROPHYSICO-TECHNICAL MACHINERY			
		PETROPHYSICO-TECHNICAL MACHIN					



SEARCHED SERIALIZED INDEXED FILED

197 AND IMP. ORDERS

SEARCHED SERIALIZED INDEXED FILED

Synthesis of new diene hydrocarbons with a conjugated system of double bonds. VI. V. I. Rauter and V. V. Moshkovova (Ural State Univ., Sverdlovsk, U.S.S.R.) 19, 1023 (1959); cf. C.A. 52, 1619. - With the improved Grignard technique (C.A. 39, 6059) involving primarily low temp., high order of diln., and slow addn. of the Grignard reagent, a no. of diene hydrocarbons were prep'd. in satisfactory yields from *n*,*d* unsat'd. ketones. Generally, ketones of higher mol. wt. give lower yields, and Me ketones react less readily than *i*Pr ketones, as far as the normal reaction is concerned. Aromatic Grignard reagents favor abnormal course of the reaction to a greater extent than Me or Et Grignard reagents, in contradiction to Köller (C.A. 2, 424). *BuMgI* (from 3.8 g. Mg and 20 g. EtI) in 40 cc. *Bu*₂O was added over 6 hrs. at -14° to 20 g. 3-methyl-3,5-heptadien-5-one in 100 cc. *Bu*₂O. After working up as described in C.A. 39, 30389, 52.9% 3-methyl-5-ethyl-3,5-heptadiene, b.p. 154°, d₂₅²⁰ 0.7710, n_D²⁰ 1.4456 was obtained. To 40 g. 4-octen-3-one, b.p. 171-3°, d₂₅²⁰ 0.8541, n_D²⁰ 1.4653, in 150 cc. *Bu*₂O, cooled to -15°, *ReMgI* (from 11.6 g. Mg and 60 g. EtI) in 80 cc. *Bu*₂O was added over 6 hrs.; after standing overnight and decompsn. with ice water, the dried *Bu*₂O ext., treated with hydroquinone, distilled, in CO₂ over 2 g. KHSO₄ and 5 g. anhyd. Na₂SO₄, and finally purified by repeated distns. over Na, gave 26.6% 3-ethyl-3,4-octadiene, b.p. 167°, d₂₅²⁰ 0.7727, n_D²⁰ 1.4626; the hydrocarbon (3 g.) and 3.2 g. maleic anhydride, heated in 10 cc. Melts in a sealed tube to 100° for 20 hrs., gave an adduct forming an Ag salt with 34.74% Ag, which appears to correspond most closely to an adduct of the hydrocarbon dimer; inspection of the condensation at room temp. with 10 min. heating to 60° gave an only product contg. 37.71% Ag. Briefly, the olefin (2 g.) heated with 2.4 g. quinone in 10 cc. Melts to 100° for hrs., gave 0.9 g. *C₁₂H₁₆* (m.p. 332°), which readily reacts with 1.6 g. *C₆H₅COCl*. To 10.5 g. *PtCl₆CHCOMe*, b.p. 104-7°, d₂₅²⁰ 0.8616, n_D²⁰ 1.4447, in 100 cc. *Bu*₂O cooled to -10° to -15°, *Strengl* (from 2.3 g. Mg and 17 g. Melts) in 30 cc. *Bu*₂O was added over 6 hrs.; treatment as above gave 16% 2-methyl-1,3-diene-5-one, b.p. 127-8°, d₂₅²⁰ 0.7452, n_D²⁰ 1.4429, while a doubled amt. of *MnMgI* gave 29.5% 1,4-diol, while a doubled amt. of *MnMgI* gave 29.5%; the Dieck-Alder adduct with maleic anhydride gave a 14% salt, contg. 33.95% Ag (corresponding to the adduct of the dimer or trimer). To 17.0 g. *PtCl₆CHCOMe* in 100 cc. *Bu*₂O, cooled to -10° to -15°, *PhMgI* (from 6 g. Mg and 38 g. PhI) in 80 cc. *Bu*₂O was added over 6 hrs. to yield, upon dehydration as above, but at reduced pressure, and vacuum distn. over Na, and finally atm. distn. over Na, 2.2 g. 2-methyl-1,3-diene-5-one, b.p. 234-7°, d₂₅²⁰ 0.8556, n_D²⁰ 1.4520; the maleic anhydride adduct (prepd. at 120°) gave a Ag salt, contg. 40.97% Ag, which was fairly close to *C₁₂H₁₆O₂*.

Aluminum chloride-catalyzed addition of *tert*-butyl chloride to propylene. Verle A. Müller (General Motors Corp., Detroit, Mich.). J. Am. Chem. Soc. 69, 1704-8 (1947). - When 1.23 moles of *C₃H₆* is bubbled into 1 mole *tert*-BuCl (1) and 2 g. AlCl₃ at -10°, at such a rate that the absorption of the gas is complete, 8-10% of the *C₃H₆* (and 23-35% 1 are recovered and 70% of a butyl chloride


A19.114 METALLURGICAL LITERATURE CLASSIFICATION

CROSS REFERENCES

SEARCHED SEARCHED SERIALIZED INDEXED SEARCHED SERIALIZED INDEXED SEARCHED SERIALIZED INDEXED FILED

fraction (II) is formed. A curve shows the effect of reaction temp. and quantity of catalyst. About 45% of II is composed of 2-chloro-4,4-dimethylpentane; the remainder is principally 3-chloro-2,3-dimethylpentane with a small quantity of the 2-Cl isomer. These were identified by reaction with the Grignard reagent and comparison of the phys. properties and derivs. with known compds. The principal fraction of the higher-boiling mixt. obtained is composed chiefly of dicyl chlorides which were not further identified. At higher reaction temps. and when more catalyst was used, the reaction was more complex; the amt. of II decreased and the primary addn. product was converted almost entirely into the rearranged product; the quantity of higher-boiling product obtained increased; the quantity of lower-boiling side-reaction products increased from a negligible to a significant quantity. 2,3-Dimethyl-3-pentanol yields a *pentaether*, m. 48.5-9°; that from 4,4-dimethyl-3-pentanol m. 52-3°.

C. J. West



ESSAFOV, V. I.

"On the Attempt to Synthetize the Ketones of the Cyclopentene Series. II"
(p. 1518)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii), 1947, Vol. 17, No. 8

PA 64/49T19

YESAFOR, V. L.

USSR/Chemistry - Diene Synthesis Jun 49
Chemistry - Grignard Reaction

Preparating and Isolating New Diene Hydrocarbons With Conjugate Double Bonds, T. I. Yesafor, Lab of Org Chem, Ural State U, Sverdlovsk, 14 PP

Zhur Obshch Khim* Vol XIX, No 6 - P.1063

Gives synthesis of 2, 6-dimethylheptadiene-1, 5; 3-methyloctadiene-2, 4; 2-tertiary-butyl-, 2, 5-dimethylhexadiene-1, 3; and 2, 3, 4-trimethylhexaene-1, 3. Physicochemical constants of 2, 6-dimethyloctadiene-4, 6 are also more precise, and yield of 2, 4-.

64/49T19

USSR/Chemistry - Diene Synthesis Jun 49
(Contd)

Dimethylpentadiene-1,3 is improved. Studies Grignard reaction of three following systems: (1) 3-methylheptene-3-on-5 and CH_3MgI , (2) 4-dimethylhexane-3-on-5 and $\text{C}_2\text{H}_5\text{MgBr}$, and (3) butyldiene-diethylpropylene ketone and $\text{C}_2\text{H}_5\text{MgBr}$. Describes tertiary alcohols, 2, 6-methyloctane-4-on-6 and 3, 4, 5-trimethylheptene-4-on-3. Shows that in the dehydration of α, β -unsaturated tertiary alcohols, having in the presence of carbinol the groups CH_3 and C_2H_5 , activity is chiefly caused by the H-stem of ethyl. Submitted 9 Sep 47.

64/49T19

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0

CA

The characteristics of α,β -unsaturated ketones. IV.
V. I. Eslov. J. Gen. Chem. U.S.S.R. 19, 1100-18
(1949) (Engl. translation).—See C.A. 44, 5795a. E. J. C.

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0"

CH

Characterization of α,β -unsaturated ketones. IV. V. I. Basov, Zhur. Obschch. Khim. (J. Gen. Chem.) 19, 1118-24 (1949); cf. C.A. 39, 9181. — Summarization of the previous data (C.A. 39, 3958; 40, 6405; 44, 10016) shows that the degree of conjugation of C:C and C:O links decreases the reactivity of unsatd. ketones; increased mass of R bound to the CO group decreases the extent of conjugation, and the same takes place with the R attached to the terminal C of the C:C link. Two satd. radicals on the latter C atom decrease the conjugation even more, and the action of a t -Me-Et combination is greater than that of di-Me or t -Me and *tert*-Bu. Introduction of a satd. R on the 2nd C atom, with 2 radicals on the 1st C atom, gives the max. lowering of the conjugation. The iodination of the unsatd. ketones may serve to show the structural types by increase of the iodine no. with increased no. of side chains from the C atoms of the conjugated

system; bromination in CCl_4 can also be used similarly, as the iodine no. of addn. decreases with an increased no. of radicals bound to the conjugated system. The upper conjugation limit is given by vinyl ketones and the lower limit lies with the compds. having 2 quaternary C atoms in the C:C link. The ketones used (the methods of prepn. will be given in later paper) were: *butylideneacetone* (I), b_{10} 163-8°, d_4^{20} 0.8520, n_D^20 1.4124; *butylideneethyl ketone* (II), b_{10} 175-7°, d_4^{20} 0.8413, n_D^20 1.4470; *butylidene-methyl propyl ketone* (III), b_{10} 183-7°, d_4^{20} 0.8540, n_D^20 1.4450 (probably contains some 3-methyl-3-hepten-2-one); *enanthylideneacetone* (IV), b_{10} 100-10°, d_4^{20} 0.8473, n_D^20 1.4480; *isooamylideneacetone* (V), b_{10} 174-8°, d_4^{20} 0.8470, n_D^20 1.4400; *meityl oxide* (VI), b_{10} 129-30°, d_4^{20} 0.8602, n_D^20 1.4446; *3-methyl-3-hepten-5-one* (VII), b_{10} 164°, d_4^{20} 0.8591, n_D^20 1.4421; *3,3-dimethyl-3-hexen-5-one* (VIII), b_{10} 157-8°, d_4^{20} 0.8585, n_D^20 1.4415; *3-ethyl-4-methyl-3-hepten-2-one* (IX), b_{10} 107-91°, d_4^{20} 0.8543, n_D^20 1.4450; *2,2,5,5,6,6-pentaethyl-4-hepten-3-one* (X), b_{10} 202-3°, d_4^{20} 0.8525, n_D^20 1.4401; *benzylideneacetone* (XI), m 42.5°; and *beetylideneethyl ethyl ketone* (XII), m 40°. The iodination was run by addn. of the ketone to 10 ml. EtOH, rinsing with 10 ml. EtOH, addn. of 25 ml. EtOH-iodine soln. and, after shaking, 200 ml. H_2O .

followed by shaking and letting stand 5 min. or longer, and titration of the excess iodine with thiosulfate, while III is calcd. from the reaction with 10 ml. 2% KIO_3 and a 2nd titration with thiosulfate; brominations were done in CCl_4 , the mixts. allowed to stand 30 min. after mixing, and the HBr generated calcd. similarly from the reaction with KI and KIO_3 . In 5 min. I gave an iodine no. of 3.6-8.9 with 6.9-6.1% III formation, while a 1370-min. reaction gave 37.6-32.2 and 79.1-83.5%, resp.; II in 5 min. gave 6.4-6.1 iodine no. and 20.1-32.4% III, while 1330 min. gave 25.7-19.7 and 67.3-78.5%, resp.; III in 5 min. gave 24.5-5.2 and 10.5-18.0%, resp. (longer runs not made); XI in 5 min. gave 5.1-3.0 iodine no. (III not reported) while 1350 min. gave 15.1-18.6 and 15.8-17.0%, resp.; XII in 5 min. gave 8.8-5.9 and 36.6-7.0%, resp.; VII gave in 5 min. 3.2-6.5 iodine no. and no III cited, while 1740 min. gave 105.4 and 40.0% III, resp.; X in 5 min. gave 4.8-6.1 iodine no. and 33.1-44.0% III; VII in 5 min. gave 36.6-4.5 iodine no. and 93.6-8.4% III, while 1700 min. gave 150.7-3.5 and 59.2-64.0%, resp.; VIII in 5 min. gave 90.3-89.2 and 69.0-5.4%, resp., while in 1410 min. it gave 198.8-204.4 and 80.1-79.0%, resp.; IX in 5 min. gave 109.7-5.7 and 82.6-72.3%, resp.; IV in 5 min. gave 30.8-8.3 and 40.2-2.0%, resp. The bromination expts. gave the following nos. of substitution and addn., resp.: I 17-32 and 194-6; II 21-7 and 185-90; III 49-7 and 157-61; IV 17-14 and 162-55; XI — and 161; XII 6-9 and 135-6; VI 11-10 and 249-6; X 23-19 and 114-16; VII 179-87 and 119-14; VIII 312-36 and 19-18; IX 223-13 and 50-64. The yields of Grignard reactions with the above ketones, cited from previous papers (references given above), are also affected by the extent of conjugation, but the results must be interpreted with care since 1,4-addn. is also possible, although the generally higher yields are assumed, with systems having a

low degree of conjugation, with allowance being made for the ease of enolization. V. V. I. Riafov and V. V. Sergovskaya. *Ibid.* 17(2) 3.—Lithiidesoacetonone (I) has the highest degree of conjugation of unsat. in the RCH_2CHAc series, i.e., increase of R increases the independence of the ethylene bond. It does not react in aq. alc. soln. with 0.1 N iodine, while with 0.2 N soln. the iodine no. is but 1.57% of theory, and even a 0.6 N soln. gives but 2.30% iodine no. (of theory); the butyliidene analog gives 1.53, 4.39, and 17.75% of the theoretical iodine no. under the same conditions. Best prepn. of I: 650 ml. Me_2CO and 700 ml. 1.4% aq. NaOH treated at 0° with 120 g. AcH in 280 ml. H_2O , precooled to -2° over 6 hrs., let stand overnight at 0°, neutralized with AcOH, and satd. with NaCl, gave on evap. with PtO_2 and distn. in the presence of a crystal of iodine, 18.6% of the product, $b.p.$ 119-21°, $d^{20}_{4} 0.8814$, $n_D^{20} 1.4339$. Bromination of I at 0° in CCl_4 gives 88% of the theoretical uptake, even with twice the theoretical amt. of Br; no HBr is evolved. G. M. Kosolopoff

2/50T30

YESAFOV, V. I.

USSR/Chemistry - Alcohols
Synthesis

Jul 49

"The Synthesis of Tertiary Alcohols Using Camphor
as a Base," I., V. I. Yesafov, N. I. Novikov,
Lab of Org Chem, Ural State U, Sverdlovsk, 62 PP

"Zhur Obshch Khim" Vol XIV, No 7

Yield of tertiary methylborneol, synthesized by
interaction of camphor and CH₃Li, is decreased
at high temperatures chiefly because of enoliza-
tion of camphor. At 150°, emulsionization of camphor
is accompanied by formation of secondary compone-
nts containing camphor, latter reaction being chief
cause of a reduced yield of tertiary alcohol

2/50T30

USSR/Chemistry - Alcohols
Synthesis (Contd)

Jul 49

When synthesis is carried out at -150° C. At high
temperatures chief secondary reaction is reduc-
tion of camphor into borneol. Submitted
29 Mar 48.

2/50T30

CA

Characterization of α,β -unsaturated ketones. VI.
V. I. Esafoy (Ural State Univ., Sverdlovsk). *Zhur. Obrabotki Khim.* (J. Gen. Chem.) 20, 1050-7 (1950); cf. C.I. 44, 6810d.— α,β -Unsatd. ketones with normal structure are more readily hydrolyzed with 0.01 N than with 0.1 N NaOH; ketones similar to mesityl oxide behave in the reverse manner. Such gradation of reactivity may be used in the characterization of ketones. Ketones analogous to trialkylvinyl ketone, i.e. $R_3C:CCOR$, are even more resistant to hydrolysis. In the 1st series 0.1-0.2-g. samples were heated in ampuls with H_2O , 0.01 or 0.1 N NaOH (10 ml.) 2 hrs. at 100°, and the CH_4 reaction on 10-ml. aliquots used as the relative measure of iodine consumption, calcd. on the starting material. In a 2nd series the alk. iodine soln. was allowed to react with the 10-ml. aliquot for 15 sec., 1, 5, or 15 min. so as to give a measure of the abs. amts. of the reacted ketones. The results follow. PyCH:CHCOMe: in H_2O 01.33% reacted, in 0.01 N NaOH 80.91%, in 0.1 N 73.5%. $CH_3CH:CHCOMe$, 60.3, 81.0, 7171%, resp. Mesityl oxide, 44.68, 75.43, 90.78%. EtCOMe:CHCOEt, 43.23, 60.16, 92.10%. EtCOMe:CMeCOMe, 37.08, 37.49, 46.89%. The ketones with normal structure have a higher degree of conjugation than occurs in branched-chain analogs and this is held to be the reason for their more easy cleavage.

C. W. Wilson

CA

7

Use of the Iodoform reaction in the analysis of some er-
gic compounds...V. I. Ralov and N. M. Stafeva (A. M.
Gor'kiy Ural State Univ., Sverdlovsk, U.S.S.R.). *Zhur.*
Anal. Khim., 6, 195-200 (1961). -The iodoform reaction,
i.e., the reaction of compds. contg. the group $\text{CH}_3\text{CO}-$ or
 $\text{CH}_3\text{CH}(\text{OH})-$ with I followed by hydrolysis and forma-
tion of CHI_3 was pos. also with α,β -unsatd. ketones and
their corresponding ketols not having these groups but cap-
able of yielding acetaldehyde or satd. methyl ketones upon
hydrolysis, e.g., 3-methyl-3-heptenes-3-one and 4-octene-3-
one. Therefore, the data based on CHI_3 reaction of α,β -
unsatd. methylethenes and of their β -ketols which upon
hydrolysis yield 2 mols. of satd. methyl ketone or aldehyde
and methyl ketone is inaccurate. M. Hersh

USSR/Chemistry - Diene Hydrocarbons

Apr 52

"Bromination and Iodoxydation of 1,1-Dialkylbutadiene-1,3." Part VIII," V. I. Yesafor, Lab of Org Chem., Urul State U

"Khimi Obshch Khim" Vol XXII, No 4, pp 604-611

Debromination or dialkyl allyl carbinols in presence of iodine proceeds in 2 directions with the formation of diene hydrocarbons both with isolated bonds. The yield of the latter is the lower, the lower the degree of hydrogenation of the allyl group attached to the phenol C-atoms. In 1,1-dimethylbutadiene-1,3,

224T36

contrast, to di-isopropenyl and hexadiene-2,4, the double bonds are less strongly conjugated, and the bromination or iodoxydation proceeds more fully. Tertiary sites with an allyl radical split at the bond between the radical and the C-atom of the carboxyl radical.

YESAFOV, V.I.

224T36

YESAFOV, V.I.

Role of Russian chemists in developing methods of synthesis by
means of zinc organic compounds. Trudy Inst. int. est. i tekhn. vol.6:
318-346 '55. (MLIA 9:5)
(Zinc organic compounds)

"APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0

RECORDED BY: [REDACTED] ON: [REDACTED]

APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0"

"APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0

APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0"

"APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0

APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0"

"APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0

APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962920002-0"

YESAFOV, V.I.

Chemical structure of C₁₂H₁₄ and synthesis of new phenylated
diene (1,3) hydrocarbons. Part 7. Zhur. ob. khim. 27 no.10:2667-2675
O '57. (MIRA 11:4)

1.Ural'skiy gosudarstvennyy universitet.
(Chemical structure) (Hydrocarbons)

AUTHOR: Yesafov, V. I. 79-2d-4-17/69

TITLE: Investigation of the Thermal Decomposition of Magnesium Iodide Ethers (Исследование термического разложения магниево-иодидных эфиров) I. On the Chemistry of Oxonium Compounds (I. Khimii oksoniyevykh soyedineniy)

PERIODICAL: Zhurnal Otechestvennoi Khimii, 1958, Vol. 28, Nr 5, pp. 1212-1218 (USSR)

ABSTRACT: The present investigation aimed at determining the degree of the binding stability of the ether molecules of the mentioned compounds; for this also the method of thermal decomposition was used which has to be preferred in the given case to other purely chemical methods. It was found that the ethers form diethers with magnesium iodide and this with a weak binding of the magnesium with the oxygen of the ether molecules whereby they decompose into thermal decomposition to magnesium iodide and ether. It showed that the ethers with regard to their capability to form various kinds of magnesium iodide diethers differ from each other. Some ethers form one and the same magnesium iodide

Card 1/5

Investigation of the Thermal Decomposition of Magnesium Iodide Ethers.

I. On the Chemistry of Oxonium Compounds

disodium, even though it was formed from magnesium iodide decomposition of a mixture of magnesium and iodine. The second ether group yields magnesium iodide diethers of two kinds. The first is obtained of magnesium iodide as such, the other from a mixture of iodine and magnesium or by displacement from the ethyl ether from $MgI_2 \cdot 2(C_2H_5)_2O$. Ethers with light primary radicals are displaced from $MgI_2 \cdot 2O$ by ethers with heavy primary radicals. The thermal decomposition of the magnesium iodide diethers of the second kind with aliphatic ethers is accompanied by a formation of olefins and alkyl iodides with one ether molecule and iodine atom taking part in the chemical reactions. The second iodine atom remains strongly bound to the magnesium. It was further shown that in the thermal reaction of the magnesium iodide-ethers of the second kind with ~~fat~~ aromatic ethers specific alkyls phenols and resinous masses form. One molecule of the ~~fat~~ aromatic ether separates in free state. The breaking of a solid chemical bond of magnesium to the oxygen

Card 2/3

Investigation of the Thermal Decomposition of Magnesium Iodide Ethers. 79-00-1-37/65

I. On the Chemistry of Oxonium Compounds

of the ~~the~~ ether molecule in the magnesium iodide diether of the second kind proves its oxonium nature. From a scheme for the thermal decomposition of the magnesium iodide diether of the second kind was proposed. There are 6 papers, 5 of which are Soviet.

ASSOCIATION: Uralskiy gosudarstvennyy universitet
(Ural State University)

SUBMITTED: May 6, 1967

Card 3/4

AUTHOR: Yesafov, V. I. 79-28-5-18/69

TITLE: Investigation of the Thermal Decomposition of the Reaction Products of Dioxane With Magnesium Iodide-Diether , Magnesium-Iodide and the Mixture of Magnesium and Iodine (Izuchenie termicheskogo razlozheniya produktov vzaimodeystviya dioxana s diefiratom yodistogo magniya, yodistym magniyem i smes'yu magniya s yodom)II. On the Chemistry of Oxonium Compounds (II. K khimii oksoniyevykh soysedineniy)

PERIODICAL: Zhurnal Obshchey Khimii, 1958, Vol. 28, Nr 5, pp. 1218 - 1221 (USSR)

ABSTRACT: After it was found that ethers with magnesium iodide yield compounds of the formula $MgJ_2 \cdot 2ROR'$ which on heating decompose again to their initial products, it was of interest to investigate analogous compounds containing dioxane. Above all it was found that in the action of dioxane on $MgJ_2 \cdot 2(C_2H_5)_2O$ and $RMgHal \cdot 2(C_2H_5)_2O$ (Reference 2) a complete analogy exists. The dioxane displaces equimolecularily the ethyl ether from these compounds. Different from the ethers, the compound

Card 1/3

79-28-5-16/69

Investigation of the Thermal Decomposition of the Reaction Products of
Dioxane With Magnesium Iodide-Diether, Magnesium Iodide and the Mixture of
Magnesium and Iodine . II. On the Chemistry of Oxonium Compounds


MgJ₂ with dioxane, MgJ₂.2C₄H₈O₂, on heating separates only half of the dioxane contained in it, the other half suffering decomposition. Thus the oxygen atoms of dioxane have a greater capability to form oxonium salts than the ethers, and they yield more stable oxonium compounds. The most typical oxonium compound of dioxane is obtained in its reaction with a mixture of magnesium and iodine. This product must be regarded as oxonium salt of symmetrical (formula I) or asymmetrical structure (II). The question if these salts form at the same time, or if first salt (I) is formed which then converts to salt (II) remains unsettled. It is only clear that in salt (I) if it remains unchanged on heating the β,β'-diiodine-diether would have to be found among the decomposition products, which was not observed. The experimental results speak fully in favor of structure (II). Among the decomposition products of this salt there were found: dioxane, ethylene, ethylene iodide.

Card 2/3

79-28-5-18/69

Investigation of the Thermal Decomposition of the Reaction Products of Dioxane With Magnesium Iodide-Diether, Magnesium Iodide and the Mixture of Magnesium and Iodine. II. On the Chemistry of Oxonium Compounds

ethyl iodide, acetic anhydride, iodine and magnesium. After all the results of the reaction of dioxane with the mixture of magnesium and iodine prove that the latter as well as the ethers in this case do not react according to the scheme



as a high-melting dioxonium salt is formed here instead of the low-melting ethylene iodide. There are 2 Soviet references.

ASSOCIATION: Ural'skiy gosudarstvenny universitet (Ural' State University)

SUBMITTED: May 3, 1957

Card 3/3

5 (3)

AUTHORS:

Yesafow, V. I., Stashkov, L. I., Sirotkin, L. B.,
Suvorov, A. L., Novikov, Ye. G.

SOV/79-29-3-20/6

TITLE:


On the Characteristics of the α,β -Unsaturated Ketones. VII
(Kharakteristika α,β -nepredel'nykh ketonov. VII)

PERIODICAL:

Zhurnal obshchey khimii, 1959, Vol 29, Nr 3, pp 845-849 (USSR)

ABSTRACT:

The present paper is issued as first publication of experimental data on the hydrolytic cleavage of the aliphatic aromatic α,β -unsaturated ketones containing an aryl radical which is directly combined with the carbonyl group. Ketones of this type were obtained by dehydration of the β -ketols which had been synthesized according to the method of Grignard, V. and Colonge, I. (Ref 2).



Experiments with respect to the hydrolytic cleavage of the β -ketols were carried out as well. The data of table 2 show that the β -ketols are far more unstable than the corresponding

Card 1/2

On the Characteristics of the α,β -Unsaturated Ketones. VII

SOV/79-29-3-20/61

α,β -unsaturated ketones and prove to be more sensitive to very weak hydrolysis reagents. Besides, the behavior of the β -ketols in the hydrolysis differs from that of the α,β -unsaturated ketones by the fact that a change of the NaOH-concentration exerts a slight influence upon the cleavage intensity of the β -ketols whereas the hydrolytic cleavage of the α,β -unsaturated ketones is considerably influenced. The rate of hydrolysis of the aliphatic aromatic ketones investigated increases significantly when the NaOH concentration is increased from 0.01 to 0.1 n. 8 β -ketols hitherto unknown were synthesized and described. It was determined how far the hydrolytic cleavage of the β -ketols and at the same time that of the α,β -unsaturated ketones develops and it was proved that the latter separate but little HBr on bromination. There are 2 tables and 4 references, 2 of which are Soviet.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet (Ural State University)

SUBMITTED: February 18, 1958

Card 2/2

5(3)

AUTHOR: Yesafov, V. I.

SOV/79-29-7-5/83

TITLE: Synthesis of the Phenylated α,β -Unsaturated Alcohols and the (1,3)-Diene Hydrocarbons.VIII (Sintez fenilirovannykh α,β -nepredel'nykh spirtov i diyenovykh (1,3) uglevodorodov.VIII)

PERIODICAL: Zhurnal obshchey khimii, 1959, Vol 29, Nr 7, pp 2129-2132 (USSR)

ABSTRACT: In order to solve the problems concerning the separation of $HOMgX$ from the forming magnesium halide carbinolates (Refs 1-3), the reaction of 2,2,3-trimethyl-5-p-arylpenten-3-ones-5 ($Ar=C_6H_5$, $n=CH_3C_6H_4$, $n=C_2H_5C_6H_4$), which may be present in cis- and trans-forms with MgX (Ref 4), are described here. The experiments showed that among the three homologous ketones 2,2,3-trimethyl-5-p-tolylpenten-3-one-5 produces the best yields in tertiary alcohols. These alcohols can be dehydrated only with difficulties. In the Grignard reaction the other two ketones mainly yielded diene hydrocarbons (1,3). This particular behavior might be explained also by the influence of a spatial factor. The phenylated diene hydrocarbons (1,3) of the structure (III) and (IV) obtained show, compared to the similar aliphatic compound (V) (Ref 3), a stronger stability of the double bond

Card 1/2

Synthesis of the Phenylated α,β -Unsaturated
Alcohols and the (1,3) Diene Hydrocarbons. VIII

SOV/79-29-7-5/83

in position 3,4 and may be brominated therefore only with the separation of a high amount of HBr. The following α,β -unsaturated tertiary alcohols were newly synthesized and characterized: 2,2,3-trimethyl-5-p-tolylhexen-3-ol-5 and 2,2,3-trimethyl-5-p-tolylhepten-3-ol-5 as well as the phenylated diene hydrocarbons (1,3): 2-phenyl-4,5,5-trimethylhexadiene-1,3 and 2-p-ethyl-phenyl-4,5,5-trimethylhexadiene-1,3. It is assumed that the stability of the halogen magnesium carbinolates formed by Grignard reaction depends on α,β -unsaturated ketones. There are 1 table and 4 Soviet references.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet (Ural State University)

SUBMITTED: March 17, 1958

Card 2/2

YESAFOV, V. I.

On the history of the origin of the stereochemical theory and
the attitude of A.M. Butler and some of his contemporary West
European chemists toward it. Trudy Inst.ist.est.i tekhn. 304-6.
135-174 '60. (MIRA 13:8)

(Stereochemistry)
(Butlerov, Aleksandr Mikhailovich, 1828-1886)

86499

53610 2209, 1373, 1153

S/079/60/030/011/004/026
B001/B066

AUTHORS: Yesafov, V. I. and Yakunina, G. I.

TITLE: Chemistry of Onium Compounds. III. Investigation of Thermal Decomposition of the Reaction Products of Tetrahydrofuran, α -Methyl Furan, Pyrrole, Thiophene With the Dietherate of Magnesium Iodide and With Magnesium Iodide


PERIODICAL: Zhurnal obshchey khimii, 1960, Vol. 30, No. 11,
pp. 3572-3576

TEXT: V. I. Yesafov (Refs. 1,2) showed in his papers that the etherate $MgI_2 \cdot 2(C_2H_5)_2O$ is a very convenient agent for the relative estimation of the degree of aromaticity of the five-membered O-, N- and S-heterocyclic compounds. These heterocyclic compounds may be arranged in the following order on the basis of increasing difficulty in the release of the unshared electron pair of heteroatoms with formation of "onium compounds"; and for silvan, pyrrole, and thiophene with respect to the degree of stability increase of the electron sextet of heterocycles:

Card 1/4

86499

Chemistry of Onium Compounds. III. Investigation S/079/60/030/011/004/026
of Thermal Decomposition of the Reaction B001/B066
Products of Tetrahydrofuran, α -Methyl Furan, Pyrrole, Thiophene With the
Dietherate of Magnesium Iodide and With Magnesium Iodide



This order is further confirmed by the experimental data of thermal decomposition of the reaction products of the mentioned heterocyclic compounds with anhydrous magnesium iodide. Tetrahydrofuran forms with the latter a compound which decomposes on heating by cleaving the heterocycle (Ref. 5). Silvan and pyrrole give, only on heating with magnesium iodide, compounds which decompose at high temperature, also under cleavage of the heterocycles. Also on prolonged heating, thiophene does not react with magnesium iodide. It follows from this that furan and pyrrole, as well as their compounds, maintain the "benzoid-like" state of electrons in the heterocycles only at low temperatures, with increasing temperature, however, this state is disturbed, in which connection the unshared electron pairs of oxygen and nitrogen are set free, and stable onium compounds are formed with MgI:

Card 2/4

86499

Chemistry of Onium Compounds, III. Investigation 8/079/60/030/011/001/026
 of Thermal Decomposition of the Reaction 8001/3066
 Products of Tetrahydrofuran, α -Methyl Furan, Pyrrole, Thiophene With the
 Dietherate of Magnesium Iodide and With Magnesium Iodide



which also explains the decomposition of C- and N-heterocycles on heating. It was thus shown that 1) tetrahydrofuran displaces the diethyl ether from $MgI_2 \cdot 2(O_2H_5)_2O$ to form $MgI_2 \cdot 2O \cdot H_2O$ which is decomposed under cleavage of a molecule of tetrahydrofuran, that 2) silvan and pyrrole displace one molecule of ether from $MgI_2 \cdot 2(O_2H_5)_2O$, and give compounds with MgI_2 which decompose under cleavage of the heterocycles, that 3) thiophene reacts neither with $MgI_2 \cdot 2(O_2H_5)_2O$ nor with MgI_2 . It is assumed that the participation of heteroatoms in the formation of heterocycles increases their capability of forming onium compounds. This assumption is supported by the fact that dioxane and tetrahydrofuran give with MgI_2 stabler compounds than simple aliphatic ethers. G. I. Kusnetsova, I. T. Bel'skiy,
 Card 3/4

86439

Chemistry of Onium Compounds. III. Investigation 6/079/60/030/011/004/026
of Thermal Decomposition of the Reaction Products of Tetrahydrofuran, α -Methyl Furan, Pyrrole, Thiophene With the Dietherate of Magnesium Iodide and With Magnesium Iodide

and S. Z. Tayts are thanked for making available the samples. There are 11 references: 5 Soviet, 4 US, 2 British, and 2 German.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet (Ural State University)

SUBMITTED: August 18, 1959

Card 4/4

YESAFOV, V.I.

Characteristics of diisotertiary β -glycols. Part 1. Zhur. ob. khim.
30 no.10:3272-3275 O '61. (MIRA 14:4)

1. Ural'skiy gosudarstvennyy universitet.
(Glycols)

YESAFOV, V. I.

39276

S/190/62/004/006/002/026
E101/B110

15. 8220

AUTHORS: Tager, A. A., Suvorova, A. I., Goldyrev, L. N., Yesafov,
V. I., Berestova, V. L.

TITLE: Effect of the chemical structure of the plasticizer on the
vitrification temperature of polymers. I. Plasticizing of
polystyrene with diphenic acid and naphthalic acid esters

PERIODICAL: Vysokomolekulyarnyye soyedineniya, v. 4, no. 6, 1962,
803-808

TEXT: Thermomechanical curves were plotted for polystyrene (PSt)
plasticized with 25 mole% of: monomethyl-, monoethyl-, and monobutyl
diphenate; dimethyl-, diethyl-, ethyl-butyl-, dibutyl-, ethyl-octyl-, and
diheptyl diphenate; dimethyl, diethyl, and dibutyl naphthalate. The
synthesis of ethyl-butyl diphenate (b.p. 167-168°C/15 mm Hg, MR 91.89)
and of ethyl-octyl diphenate (MR 110.57), now produced for the first time,
will be published. The compatibility of the plasticizer with PSt was
studied on the basis of the critical mixing temperature, which lay at
100-130°C with diphenic acid monoester, below room temperature (sometimes

Card 1/2

S/190/62/004/006/002/026

B101/B110

Effect of the chemical structure ...

at $\sim -50^{\circ}\text{C}$) with esters of this acid, and at room temperature with naphthalates. Results: (1) The vitrification temperature, T_v , of plasticized PSt drops with increasing compatibility. Pure PSt had $T_v = 105^{\circ}\text{C}$, PSt with monoesters had $T_v = 40-70^{\circ}\text{C}$, PSt with diphenic acid diesters yielded the lowest T_v . T_v dropped with increasing length of the alkyl radical: ethyl-octyl diphenate yielded $T_v = -11^{\circ}\text{C}$; the naphthenates showed a low effect ($T_v = 9-48^{\circ}\text{C}$). (2) With increasing content of CH_2 links in the alkyl radical, T_v of diphenic acid diesters approaches a minimum at $n_{\text{CH}_2} = 10-12$, and then rises again. (3) The structure of the aromatic radical of the plasticizer affects T_v : diphenates (and phthalates) plasticize more intensively than naphthalates. There are 3 figures and 2 tables.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet im. A. M. Gor'kogo
(Ural State University imeni A. M. Gor'kiy)

SUBMITTED: March 21, 1961

Card 2/2

YESAFOV, V.I.

38277

S/190/62/004/C06/003/026
B101/B110

15.2.2.20

AUTHORS: Tager, A. A., Suvorova, A. I., Goldyrev, L. N., Yesafov, V. I.,
Topina, L. P.

TITLE: Effect of the chemical structure and the size of the plasticizer molecule on the vitrification temperature of polymers. II. Plasticizing of polymethyl methacrylate with esters of diphenic and naphthalic acids

PERIODICAL: Vysokomolekulyarnyye soyedineniya, v. 4, no. 6, 1962, 809-814

TEXT: Thermomechanical curves were plotted for polymethyl methacrylate (PMMA) plasticized with 25 mole% of: monomethyl, monoethyl, and monobutyl diphenate; dimethyl, diethyl, ethyl-butyl, dibutyl, and diheptyl diphenate; dimethyl, diethyl, and dibutyl naphthalate. Results: (1) The better the compatibility between polymer and plasticizer, the greater the drop in the vitrification temperature, T_v , of pure PMMA ($T_v = 100^\circ\text{C}$).

(2) T_v dropped with increasing length of the alkyl radicals of the diphenates down to a minimum (-9°C). (3) Monesters of diphenic acid and naphthalates showed a lower plasticizing effect ($T_v \sim 50^\circ\text{C}$). (4) The

Card 1/2

S/19C/62/004/006/003/026
B101/B110

Effect of the chemical structure ...

structure of the aromatic radical affects the plasticizing effect. The better plasticizing of diphenates is explained by the ability of the compound to be turned round the C-C bond between the two benzene rings. In the case of monoesters, the free COOH reduces the compatibility. (5) The molar concentration rule does not apply to the polymer plasticizer systems investigated. There are 5 figures and 1 table.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet im. A. M. Gor'kogo
(Ural State University imeni A. M. Gor'kiy)

SUBMITTED: March 21, 1961

Card 2/2

YESAFOV, V.I.

History of the discovery of organic reactions in the presence
of anhydrous aluminum halides. Trudy Inst.ist.est.i tekhn.
39:104-140 '62. (MIRA 16:2)
(Friedel-Crafts reaction)

YESAFOV, V.I.; SHITOV, G.P.

Characteristics of primary-tertiary β -glycols. Part 3. Zhur. ob.-
khim. 32 no.9:2819-2822 S '62. (MIRA 15:9)

1. Ural'skiy gosudarstvennyy universitet.
(Glycols)

YESAFOV, V.I.; ZHUKOVA, L.P.

Characteristics of secondary-tertiary β -glycols. Part 2.
Zhur. ob. khim. 32 no. 9:2816-2819 S '62. (MIRA 15:9)

1. Ural'skiy gosudarstvennyy universitet.
(Glycols)

YESAFOV, V.I.; DASHKO, V.N.; MAREK, E.M.

Characteristics of secondary-tertiary β -glycols. Part 5. Zhar.
b. khim. 34 no.12:4094-4096 D '64 (MIRA 18:1)

1. Ural'skiy gosudarstvennyy universitet.

USSR / Human and Animal Physiology (Normal and Pathological).
Lymph Circulation.

T

Abs Jour : Ref Zhur - Biologiya, No 13, 1958, No. 60370

Author : Yesakov, A. I.

Inst : Not given

Title : Investigation of the "Automatic" Action of the Center
of Lymphatic Hearts

Orig Pub : Zh. obshch. biologii, 1957, 18, No 3, 185-193

Abstract : The contraction of the posterior lymph hearts (LH) in a frog was recorded on a kymograph. The warming of the spinal cord with a thermode on the level of the III vertebra produced an increase in the contraction rate; the destruction of the cord on that level did not interrupt the rhythmical action of LH. It was stopped only when the spinal cord was destroyed at the VII vertebra. The rhythmic activity of LH originates in the

Card 1/2

Y.SAKOV, A.I., Cand. Biol. Sci.—(disc) "Study of the ~~mechanism~~^{mechanisms} of the so-called automatic ~~activity~~^{activity} of lymphatic hearts." Ios, 1958.
14 pp (Mos Order of Lenin and Order of Labor Red Banner St. to Min. S.V.Lomonosov), 150 copies

-25-

YESAKOV, A.I.

Metabolic factors of automatism (exemplified by the cerebrospinal center of lymph hearts). Zhur. ob. biol. 20 no.1:28-34 Ja-F '59.
(MIRA 12:2)

1. Kafedra fiziologii shivotnykh Moskovskogo gosudarstvennogo universiteta im. M.V.Lomonosova.
(LYMPHATICS) (SPINAL CORD) (PHYSIOLOGICAL CHEMISTRY)

SNYAKIN, P.G., prof.; YESAKOV, A.I., kand.biolog.nauk

Muscular sense. Zdorov'e 7 no.9:9-10 S '61.
(MUSCULAR SENSE)

(MIRA 14:9)

YESAKOV, A.I.

Metabolic nature of the automatism of nerve cells as exemplified
in the spinal center of the lymph heart. Zhur. ob. biol. 22 no.2:
136-143 Mr-Ap '61. (MIRA 14:5)

1. Department of Animal Physiology, State University of Moscow.
(NERVES)

YESAKOV, A.I.

Problems of the efferent regulation of receptors; based on an example of lingual chemoreceptors. Biul.eksp. biol. i med. 51 no.3: 3-8 Mr '61. (MIRA 14:5)

1. Iz laboratorii fiziologii i patologii organov chuvstv (zav. - prof. P.G.Snyakin) Instituta normal'noy i patologicheskoy fiziologii (dir. - deystvitel'nyy chlen AMN SSSR V.V.Parin) AMN SSSR, Moskva. Predstavlena deystvitel'stva chlenom AMN SSSR P.K.Anokhinym.
(RECEPTORS (NEUROLOGY)) (STOMACH)
(TONGUE)

YESAKOV, A. I.

"Methodology of neurodynamic investigations and practical work
in the physiology of human analysors" by P. O. Makarov. Reviewed
by A. I. Esakov. Nauch. dokl. vys. shkoly; biol. nauki no.3:
208-209 '62. (MIRA 15:7)

(SENSES AND SENSATION) (PHYSIOLOGY--LABORATORY MANUALS)
(MAKAROV, P. O.)

SNYAKIN, P. G., prof.; YESAKOV, A. I., kand. biologicheskikh nauk

Our habits are stereotypes. Zdorov'e 8 no.11:4-5 II '62.
(MIRA 15:10)

(HABIT)

YESAKOV, A.I.

Effect of the sympathetic nervous system on the electric activity of tongue receptors. Trudy Inst. norm. i pat. fiziol. AMN SSSR 6:63-65 '62 (MIRA 17:1)

1. Laboratoriya fiziologii i patologii organov chuvstv (zav.-prof. P.G. Snyakin) Instituta normal'noy i patologicheskoy fiziologii AMN SSSR.

YESAKOV, A.I.; ZAYKO, N.S.

Effect of guanidine on the functional activity of taste
receptors. Fiziol. zhur. 49 no.8:984-989 Ag '63.
(MIRA 17:2)

1. From the Laboratory for Physiology and Pathology of Sense
Organs, Institute of Normal and Pathologic Physiology,
U.S.S.R. Academy of Medical Sciences, Moscow.

YESAKOV, A.I.

Reflex regulation of "spontaneous" activity of tongue chemo-receptors. Biul. eksp. biol. i med. 56 no.8:7-11 Ag '63.
(MIRA 17:7)

1. Iz laboratorii fiziologii i patologii organov chuvstv (zav. - prof. P.G. Snyakin) Instituta normal'noy i patologicheskoy fiziologii (dir. - deystvitel'nyy chlen AMN SSSR prof. V.V. Parin) AMN SSSR, Moskva. Predstavлено deystvitel'nym chlenom AMN SSSR P.K. Anokhimym.

YESAKOV, A.I.

Efferent reactions in the hypoglossal nerve. Biul. eksp. bisl. i med.
56 no.11:15-18 0 [i.e. N] '63. (MIRA 17:11)

1. Iz laboratorii fiziologii i patologii organov chuvstv (zav. - prof.
P.G. Snyakin) Instituta normal'noy i patologicheskoy fiziologii (dir.-
deystvitel'nyy chlen AMN SSSR prof. V.V. Parin) AMN SSSR, Moskva.

YESAKOV, A.I.

Electrophysiological analysis of the functional mobility and of
processes regulating the taste receptor apparatus. Trudy Inst.
norm.i pat.fiziol. AMN SSSR 7:43-44 '64. (MIRA 18:6)

I. Laboratoriya fiziologii i patologii organov chuvstv (zav. -
prof. P.G.Snyakin) Instituta normal'noy i patologicheskoy fiziologii
AMN SSSR.

YESAKOV, A.I. (Moskva)

Role of different regulation levels of receptors in the
perception process. Vest. AMN SSSR 21 no.1:62-68 '66.
(MIRA 19:1)

YESAKOV, A.I.; FILIN, V.A.

Physiological characteristics of the functioning of the taste
receptor apparatus. Fiziol. zhur. 50 no.2:169-176 F '64.

(MIRA 18:2)

1. Laboratoriya fiziologii i patologii organov chuvstv Instituta
normal'noy i patologicheskoy fiziologii AMN SSSR, Moskva.

YESAKOV, A.I.

Possible specific stimulus of the "automatic" center of the lymph heart. Biul. eksp. biol. i med. 57 no.6:19-22 Je '64.

(MIRA 18:4)

1. Kafedra fiziologii zhivotnykh Moskovskogo gosudarstvennogo universiteta imeni Lomonosova.

1 E 2111 474, 11 14

YUR'YEV, YU.K.; BAZAN, V.I.; YESAFOVA, A.N.; SELIVERSHOVA, S.N. AND CHERNYAKHOVER, S.I.

"The Catalytic Transformations of Heterocyclic Compounds" Part XVI. "Synthesis of Certain Pyridine and Quinoline Derivatives of Pyrrolidine" Zhur Obshch. Khim. 10, No. 21, 1940. Moscow Order To Lenin State University imeni M.V.Lomonosova, Laboratory of Organic Chemistry imeni Academician N. D. Zelinskij. Received 26 May 1940.

U-1612, 3 Jan. 1952

YESAKOV, I.S.; YASHCHENKO, Z.G.

Interpreting vertical electric sounding curves by the T method.
Razved. i prom. geofiz. no.30:50-54 '59. (MIRA 12:12)
(Electric prospecting)

YASHCHENKO, Z.G.; YESAKOV, I.S.

Use of electric prospecting in studying the elastic properties
of igneous rocks. Razved.i prom.geofiz. no.33:23-29 '59.
(MIRA 13:4)

(Rocks--Electric properties) (Elasticity)

YESAKOV, V.A.

M.S.Bodnarskii; obituary. Izv.AN SSSR Ser.geog. no.1:95-96
Ja-F '54. (MLRA 7:2)
(Bodnarskii, Mitrofan Stepanovich, 1870-1953)

USSR/Geography Education

Card : 1/1 Pub. 15 - 8/20

Authors : Esakov, V. A.

Title : History of the origination of geography faculties at Russian universities

Periodical : Izv. AN SSSR. Ser. geog. 4, 57 - 60, July - August 1954

Abstract : History of the establishment of geography faculties at Russian universities since the second half of the 19th century.

Institution : Acad. of Sc. USSR, Institute of Natural History and Technique

Submitted :

YESAKOV, V.A.
USSR/Geophysics - Moscow University

FD-1152

Card 1/1 Pub. 129-16/23

Author : Yesakov, V. A.

Title : From the history of geography in Moscow University (up to the establishment of the chair of geography in 1884)

Periodical : Vest. Mosk. un., Ser. fizikomat. i yest. nauk, 9, No 7, 131-136, Oct 1954

Abstract : The Petersburg Academy of Sciences, founded in 1725, established in 1739 its Geographical Department. The author lists the numerous studies and expeditions undertaken after this date.

Institution :

Submitted : May 22, 1954

YESAKOV, V.A.; SOLOV'YEV, A.I., redaktor; VOLODINA, N.I., redaktor;
ZEMSKOVA, T.A., tekhnicheskiy redaktor

[D.N. Anuchin and foundation of the Russian academic geographical school.] D.N. Anuchin i sozdanie russkoj universiteteskoi geograficheskoi shkoly. Moskva, Izd-vo Akademii nauk SSSR, 1955. 180 p.
(Anuchin, Dmitrii Nikolaevich, 1843-1923) (MLRA 8:10)

YESAKOV, V.A.

M.I. Veniukov's correspondence with N.M. Przheval'skii.
Vop. ist.est. i tekhn. no.1:207-212 '56. (MLRA 9:10)

(Veniukov, Mikhail Ivanovich, 1832-1901)
(Przheval'skii, N.M.)

3(0)

SOV/10-59-3-21/32

AUTHOR: Yesakov, V.A.

TITLE: Scientific Contacts of A. Humboldt with Russian Scientists

PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya geograficheskaya, 1959,
Nr 3, pp 124-130 (USSR)

ABSTRACT: This is a review of scientific contacts of A. Humboldt with
Russian scientists. There is 1 photograph, and 18 references,
of which 15 are Russian and 3 German.

ASSOCIATION: Institut istorii yestestvoznaniya i tekhniki AN SSSR (the
Historical Institute of the Science of Natural History and Tech-
nology, AS USSR).

Card 1/1

30

SOV/10-59-4-28/29

AUTHOR: Yesakov, V.A.

TITLE: A. Humboldt's Anniversary in the USSR

PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya geograficheskaya
1959, Nr 4, p 158 (USSR)

ABSTRACT: The article is concerned with the 100th anniversary
of the death of A.v. Humboldt in the USSR. The
following personalities and organizations partici-
pated in the commemoration of the great German scien-
tist: Academician D.I. Shcherbakov, Corresponding
Member AN SSSR (AS USSR), S.V. Kalesnik, Academician
A.A. Grigor'yev, Corresponding Member AS USSR S.V.
Obruchev, the Akademiya nauk SSSR (Academy of Sciences
USSR), the Geograficheskoye obshchestvo SSSR (Geo-
graphical Society USSR), the Glavnaya geofizicheskaya
observatoriya im. A.I. Voeveykova (Central Geophysical
Observatory imeni A.I. Voeveykova), the Astronomiches-
kiy institut imeni P.K. Shternberga (Institute of

Card 1/3

SCV/10-59-4-23/29

A. Humboldt's Anniversary in the USSR

Astronomy imeni P.K. Shternberga), the Vsesoyuznoye botanicheskoye obshchestvo (All-Union Botanical Society), the Otdeleniye geologo-geograficheskikh nauk (Section of Geological and Geographical Sciences) AS USSR, the Institut istorii yestestvoznaniya i tekhniki (Institute of History of the Natural Sciences and Technology), the Sovetskoye natsional'noye ob'yedineniye istorikov yestestvoznaniya i tekhniki (Soviet National Association of Historians of Natural Sciences and Technology), the Moskovskiy gosudarstvennyy universitet im. M.V. Lomonosova (Moscow State University imeni M.V. Lomonosov), the Moskovskoye obshchestvo ispytateley prirody (Moscow Society of Naturalists), Obshchestvo sovetsko-germanskoy druzhby i nauchnykh svyazey (Soviet-German Society for the Promotion of Friendship and Scientific Connections), the Gosudarstvennyy istoricheskiy muzey (State Historical Museum), the Gosudarstvennyy muzey

Card 2/3

SOV/10-59-4-28/29

A. Humboldt's Anniversary in the USSR

izobrazitel'nykh iskusstv (State Museum of Graphic Arts), the Muzey zemlevedeniya MGU (Geographical Museum MGU), the Gosudarstvennoye izdatel'stvo geograficheskoy literatury (State Publishing House of Geographical Publications) and the Ministerstvo svyazi SSSR (Ministry of Communications USSR). There is 1 Soviet reference.

Card 3/3

YESAKOV, V.A.; SOLOV'YEV, A.I.; FEDOSEYEV, I.A., otv. red.;

[Russian geographical explorations of European Russia and
the Urals in the 19th and the beginning of the 20th century]
Russkie geograficheskie issledovaniia Evropeiskoi Rossii i
Urala v XIX - nachale XX v. Moskva, Nauka, 1964. 177 p.
(MIRA 17:11)

NAUMOV, Guriy Vasil'yevich; FEDOSEYEV, I.A., otv. red.; YESAKOV,
V.A., red.; SOLOV'YEV, A.I., red.

[Russian geographical explorations in Siberia in the 19th
century] Russkie geograficheskie issledovaniia Sibiri v
XIX - nachale XX v. Moskva, Nauka, 1965. 146 p.
(MIRA 19:1)

YESAKOV, V.A.; PLAKHOTNIK, A.F.; ALEKSEYEV, A.I.; FEDOSEYEV, I.A.,
otv. red.; SOLOV'YEV, A.I., red.

[Russian ocean and sea studies in the 19th to the begin-
ning of the 20th century] Russkie okeanicheskie i morskie
issledovaniia v XIX-nachale XX v. Moskva, Nauka, 1964.
158 p. (MIRA 18:1)

YESAKOV, Vasiliy Petrovich; PARFENOV, Eduard Yevgen'yevich;
PROZOROV, Valentin Alekseyevich; LERNER, D.M., red.

[Automated electric drive systems with regulated semi-conductor rectifiers] Sistemy avtomatizirovannogo elektroprivoda s upravliaemymi poluprovodnikovymi vypriamiteliами.
Leningrad, 1964. 35 p. (MIRA 17:11)

YESAKOV, V.P., inzhener.

Operating conditions of electric winches on ships. Sudostroenie
23 no.3:34-35 Mr '57. (MLRA 10:5)
(Winches) (Electricity on ships)

YESAKOV, V.P., kand.tekhn.nauk

Systems for electric propelling units using steady current. Sudostroenie
25 no.2:31-34 F '59. (MIRA 12:4)
(Electricity on ships)

L 33115-66

ACC NR: AP6024083

SOURCE CODE: UR/0144/66/000/002/0235/0236

AUTHOR: Zav'yalov, A. S.; Get'man, A. A.; Molchanov, V. D.; Krasuk, N. P.;
Agranovskiy, K. Yu.; Berger, A. Ya.; Groyer, L. K.; Ytsakov, V. P.; Miller, Ye. V.;
Pyatman, K. I.; Abryutin, V. N.; Gubanov, V. V.; Oranskiy, M. I.; Yevseyev, M. Ye.;
Horkin, G. B.; Sinol'nikov, Ye. M.; Avilov-Karnaudkov, B. N.; Bogush, A. G.;
Bolyayev, I. P.; Pekker, I. I.; Chernyavskiy, F. I.

46

B

ORG: none

TITLE: O. B. Bron (on his 70th birthday)

SOURCE: IVUZ. Elektromekhanika, no. 2, 1966, 235-236

TOPIC TAGS: electric engineering personnel, circuit breaker

ABSTRACT: Osip Borisovich Bron was born in 1896 in Klintsi. In 1920, he graduated from the physics-math faculty of Khar'kov Technological Institute. He became a professor in 1930. He defended his doctor's thesis in 1940. During the second world war, he was in the navy. After demobilization in 1950, Engineer Colonel Bron went to work teaching at the Leningrad Industrial Correspondence School. He became the head of the Chair of Theoretical Bases of Electrical Technology in 1958. He is closely associated with scientific and development work, and has cooperated closely in this area with the Leningrad "Elektrosila" plant since 1946. His work has been in the areas of spark-damping and high-power circuit breakers. He has published over 140 scientific works and 19 inventions. [JPRS]

SUB CODE: 05, 09 / SUBM DATE: none

Card 1/1

0915

1647

5.3300

77034
307/62-50-12-38/43

AUTHORS: Zhuze, T. P., Zhurba, A. S., Yesakov, Ye. A.
TITLE: Brief Communications. Investigation of P-V-t-N Relation
and Phase Equilibrium in Ethylene-Cyclohexane System
PERIODICAL: Izvestiya Akademii nauk. Otdeleniye khimicheskikh nauk,
1959, Nr 12, pp 2251-2253 (USSR)

ABSTRACT: Investigation of binary systems of unsaturated gases
and paraffins, naphthenes, and aromatic hydrocarbons,
presents great practical interest in view of the high
solubility of the latter in compressed unsaturated gases.
P-V-t-N relationship of the ethylene-cyclohexane system
was studied by the authors in a modified apparatus
described by Sage and Lacey (Trans. Amer. Inst. Mining
Met. Engr's., 1940, Nr 136, p 138). Isotherms $V = f(p)t$
were traced in the range from 30 to 150° for ethylene-
cyclohexane mixtures with 20 to 85 molar % ethylene, at
pressures ranging from 10 atm to pressures somewhat
above the saturation point of each mixture. Saturation
pressures and the corresponding specific volumes were
determined from these isotherms, and dew point pressures
were established in a series of separate experiments.

Card 1/3 2

Brief Communications. Investigation of
P-V-t-N Relation and Phase Equilibrium
in Ethylene-Cyclohexane System

77094
SOV/62-59-12-38/33

Data thus obtained, served to trace isotherms $p = f(N)t$ (where N is the molar share of ethylene dissolved in cyclohexane); isotherms of the equilibrium constant for ethylene and cyclohexane, $K = f(P)t$; and isotherms of molar volumes V_M of the binary system plotted against molar share N_2 of ethylene at 50 atm. In this manner, the composition of the coexisting phases, the equilibrium constants of ethylene and cyclohexane at pressures up to 100 atm, and the molar volumes of mixtures at their saturation pressures, were determined in the temperature range from 30 to 125°. Molar volume isotherms at low temperatures were practically linear up to $N_2 = 0.65$; at higher temperatures, the molar volume increases sharply starting with $N_2 = 0.50$. There are 3 figures; and 3 references, 1 U.S., 2 Soviet. U.S. reference is: B. H. Sage, W. N. Lacey, Trans. Amer. Inst. Mining. Met Engrs., 136, 138 (1940).

Card 2/32

Inst. Geology & Processing Thermal Fuels A.S. U.S.S.R.

ZHUZE, T.P.; YUSHKEVICH, G.N.; USHAKOVA, G.S.; YESAKOV, Ye.A.

Critical parameters for oil and oil-gas systems. Neft.
khoz. 41 no.6:25-31 Je '63. (MIRA 17:6)

AFANAS'YEVA, V. B.; YESAKOVA, N. P.

Relation between the snow cover and G. J. Wangenheim's types
of circulation. Trudy GGO no.151:77-80 '64. (MIRA 17:7)

ACCESSION NR: AT4046059

S/2531/64/000/168/0182/0188

AUTHOR: Yudin, M.I.(Doctor of physico-mathematical sciences); Yesakova, N. P.; Afanas'yeva, V. B.

TITLE: Preliminary evaluation of the prognostic significance of the information obtained from meteorological satellites

SOURCE: Leningrad. Glavnaya geofizicheskaya observatoriya. Trudy*, no. 166, 1964. Voprosy* interpretatsii dannykh meteorologicheskikh sputnikov (Problems in the interpretation of data of meteorological satellites), 182-188

TOPIC TAGS: meteorology, meteorological satellite, cloud, precipitation, weather forecasting, long-range weather forecasting, snow cover, radiation balance

ABSTRACT: The objective of this paper was to develop a method for the preliminary characterization of anomalies of cloud cover, the radiation balance of the underlying surface and the limits of snow and ice cover for subsequent use of such characteristics in long-range weather forecasting. The authors establish statistical relationships between such anomalies and the characteristics of future weather (temperature and precipitation); certain direct characteristics of atmospheric circulation are also analyzed in relation to future weather. Determination of the characteristics of anomalies of the cloud cover,

Card 1/3

ACCESSION NR: AT4046059

the boundaries of the snow and ice cover and the radiation balance was done using mean 10-day values for the period September-November 1948-1957. These values were mapped, after which the parameters characterizing the fields of individual elements were determined. The method used for constructing the maps and defining the characteristics of anomalies is described briefly. The state of atmospheric circulation was described using the zonal index devised by Ye. N. Blinova, the M. I. Yudin meridional index and the A. A. Rozhdestvenskiy hydrodynamic indices. These parameters were used to supplement the 10-day means of temperature and precipitation for an analysis of these values determined for a grid of points covering much of the European SSSR. Synchronous statistical relationships were established between the 10 mentioned parameters; asynchronous prognostic relationships also were determined. The ten considered parameters were correlated with temperature and precipitation for the 10 days which followed. The computations of the correlation coefficients were performed on a "Ural-1" electronic computer. Most of the results of the computations were plotted on maps, and 66 such maps were constructed. In a considerable number of cases relationships were discovered which are characterized by quite high correlation coefficients and with a stable identical

ACCESSION NR: AT4046059

sign for the entire considered area. It was found that the selected parameters generally give more information for prediction of temperature than for prediction of precipitation. However, an absence of prognostic relationships is noted on a number of maps. The method described made it possible to establish a number of parameters of the state of the atmosphere and the underlying surface which are quite closely related to the characteristic of future weather for 10 days in advance. The greater part of the parameters apply to those elements which cannot be determined globally except by use of meteorological satellites. This emphasizes the great importance of satellite observations for long-range forecasting. Orig. art. has: 2 formulas, 5 figures and 1 table.

ASSOCIATION: Glavnaya geofizicheskaya observatoriya Leningrad (Main Geophysical Observatory)

SUBMITTED: 00

ENCL: 00

SUB CODE: ES

NO REF SOV: 006

OTHER: 000

Card 3/3

AFANAS'YEVA, V.B.; YESAKOVA, N.P.

Statistical relations between the anomalies of certain
weather characteristics. Trudy GGO no.165:105-113 '64.
(MIRA 17:9)

AFANAS'YEVA, V.B.; YESAKOVA, N.P.

Relation of the shift of the boundary of snow cover to the G.IA.
Vangengeim types of circulation in the spring period. Trudy GGO
no.168:45-48 '65.

YUDIN, M.I.; YESAKOVA, H.P.; APANAS'YIWA, V.B.

Preliminary evaluation of the prognostic significance of information returned by meteorological satellites. Trudy GGO no.166:182-188 '64.

(MTRA 17:11)

YESAKOVA, R.

Objective method for determination of the liming stage of gelatin stock. R. Gorodetskaya, M. Sheremet, M. Shakhnazarova, D. Virnik, V. Smirnova, and R. Yesakova. Myasnaya Ind. S. S. S.R. 25, No. 5, 52-4(1954). —The procedure for detg. the status of the liming of gelatin stock is based on extg. a sample and detg. extd. gelatin colorimetrically by means of the biuret reaction. Results are given for extractable gelatin in bone stock at 5-day intervals for 40 days of liming. Total extractable gelatin is detd. for various bones and other gelatin stock.

M. M. Piskur

GORODETSKAYA, R.V., kandidat khimicheskikh nauk; SHAKHNAZAROV, M.Sh.,
mladshiy nauchnyy sotrudnik; SHEREMET, M.V.; VIRNIK, D.I.;
SMIRNOVA, V.Ye.; YESAKOVA, R.

Reducing losses in gelatin production. Trudy VNIIMP no.7:108-113
'55.
(MLRA 9:8)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut myasnoy promyshlennosti (for Gorodetskaya, Shakhnazarova, Sheremet); 2. Moskovskiy zhelatinovyj zavod (for Virnik, Smirnova, Yesakova).
(Gelatin)

GORODETSKAYA, R.V., kandidat khimicheskikh nauk; SHAKHNAZAROVA, M.Sh.,
mladshiy nauchnyy sotrudnik; SHEREMET, N.V.; VIRNIK, D.I.;
SMIRNOVA, V.Ye.; YESAKOVA, R.

Methods of determining the degree of liming in gelatinous tissues.
Trudy VNIIMP no.7:114-122 '55.
(MLRA 9:8)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut myasnoy promy-
shlennosti (for Gorodetskaya, Shakhnazarov, Sheremet); 2. Moskov-
skiy zhelatinovyj zavod (for Virnik, Smirnova, Yesakova).
(Gelating)